

University of Groningen

Density Functional Study of the Ground and Excited State Potential Energy Surfaces of a Light-Driven Rotary Molecular Motor (3R,3'-R)-(P,P)-trans-1,1',2,2',3,3',4,4'-Octahydro-3,3'-dimethyl-4,4'-biphenanthrylidene

Kazaryan, Andranik; Filatov, Michael

Published in:
The Journal of Physical Chemistry A

DOI:
[10.1021/jp902389j](https://doi.org/10.1021/jp902389j)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2009

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Kazaryan, A., & Filatov, M. (2009). Density Functional Study of the Ground and Excited State Potential Energy Surfaces of a Light-Driven Rotary Molecular Motor (3R,3'-R)-(P,P)-trans-1,1',2,2',3,3',4,4'-Octahydro-3,3'-dimethyl-4,4'-biphenanthrylidene. *The Journal of Physical Chemistry A*, 113(43), 11630-11634. <https://doi.org/10.1021/jp902389j>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

A density functional study of the ground and excited
state potential energy surfaces of a light-driven rotary
molecular motor

(3R,3'R)-(P,P)-trans-1,1',2,2',3,3',4,4'-octahydro-3,3'-
dimethyl-4,4'-biphenanthrylidene.

Supporting information.

Andranik Kazaryan and Michael Filatov

June 16, 2009

Table 1: The geometry parameters of the (3R,3'R)-(P,P)-trans-1,1',2,2',3,3',4,4'-octahydro-3,3'-dimethyl-4,4'-biphenanthrylidene molecular motor (see scheme 1 for the labels) in the four conformations: (P,P)-*trans*-**1**, (M,M)-*cis*-**2**, (P,P)-*cis*-**2**, (M,M)-*trans*-**1** (denoted in the table as PPT, MMC, PPC, MMT, correspondingly), optimized with B3LYP using two different basis sets and the relative B3LYP ground state energies with respect to PP-trans conformation in eV. The experimental values [1] are shown in brackets

	6-31G*/STO-3G				6-31G*			
	PPT	MMC	PPC	MMT	PPT	MMC	PPC	MMT
α	186.6	-17.8	2.2 (-3.2)	160.1	188.6	-17.6	0.9 (-3.2)	161.1
α'	-165.6	-21.2	5.4 (5.9)	151.7	-164.0	-22.7	7.2 (5.9)	150.8
β	67.1	-46.1	54.1 (54.4)	-44.0	68.6	-48.2	56.1 (54.4)	-45.1
γ	-121.8	-41.8	-94.3 (-97.0)	-44.8	-118.9	-40.3	-95.1 (-97.0)	-44.1
γ'	-121.8	-41.8	-94.3 (-96.0)	-44.8	-118.9	-40.3	-95.1 (-96.0)	-44.1
ζ	190.5	178.3	181.6	175.8	192.3	177.4	183.1	174.8
φ	124.3	120.7	124.4	122.1	123.7	120.9	124.9	122.1
1	1.356	1.377	1.362 (1.347)	1.378	1.357	1.377	1.361 (1.347)	1.378
2	1.495	1.500	1.494	1.495	1.494	1.498	1.493	1.492
3	1.399	1.402	1.400	1.406	1.393	1.396	1.394	1.399
4	1.523	1.520	1.523	1.520	1.509	1.506	1.509	1.506
5	1.569	1.558	1.562	1.558	1.544	1.535	1.538	1.535
6	1.580	1.562	1.569	1.556	1.570	1.551	1.558	1.545
7	1.532	1.559	1.541	1.563	1.532	1.557	1.540	1.559
ΔE	0.00	0.56	-0.01	0.45	0.00	0.62	0.03	0.43


Table 2: The vertical excitation energies in the (P,P)-*trans*-**1** and (P,P)-*cis*-**2** conformations. The geometries optimized with two different basis sets^a were used. The excitation energies were calculated with the use of TD-BH&HLYP and SA-REBH&HLYP and three different basis sets^b. Deviations from the experimental data are shown. All energies are in eV.

	SA-REKS				TDDFT			
	A		B		A		B	
(P,P)- <i>trans</i> -1								
	S ₁ -S ₀	Error	S ₁ -S ₀	Error	S ₁ -S ₀	Error	S ₁ -S ₀	Error
1	4.87	0.81	4.88	0.82	4.79	0.73	4.80	0.74
2	4.47	0.41	4.50	0.44	4.27	0.21	4.30	0.24
3	4.41	0.35	4.44	0.38	4.19	0.14	4.23	0.17
Exp. ^c	4.06		4.06		4.06		4.06	
(P,P)- <i>cis</i> -2								
	S ₁ -S ₀	Error	S ₁ -S ₀	Error	S ₁ -S ₀	Error	S ₁ -S ₀	Error
1	4.65	0.44	4.65	0.44	4.63	0.42	4.63	0.42
2	4.23	0.02	4.29	0.08	4.17	-0.04	4.20	-0.01
3	4.18	-0.03	4.24	0.03	4.10	-0.11	4.14	-0.07
Exp. ^c	4.21		4.21		4.21		4.21	

^a A: REB3LYP/hybrid 6-31G*/STO-3G optimized structures; B: REB3LYP/6-31G* optimized structures.

^b 1: hybrid 6-31G*/STO-3G; 2: 6-31G*; 3: 6-311G**

^c Cited from Ref. [1].



$\alpha=(4a, 4, 4',4a')$; $\alpha'=(3, 4, 4',3')$
 $\beta=(4', 4, 4a,4b)$; $\beta'=(4, 4', 4a',4b')$
 $\gamma=(4, 4', 3',15')$; $\gamma'=(4', 4, 3,15)$
 $\zeta=(3, 4, 4',4a)$; $\varphi=(3, 4, 4')$
 $1=44'$; $2=44a$; $3=4a10a$;
 $4=10a1$; $5=12$; $6=23$; $7=34$

Scheme 1:

References

- [1] Harada, N. ; Koumura, N.; Feringa, B. L. *J. Am. Chem. Soc.* **1997**, *119*, 7256.